

Overcoming Challenges in Outpatient Drug Therapy

Katie Lacaria
 Clinical Pharmacist
 Outpatient Leukemia/Bone Marrow Transplant Program of BC
 Clinical Services Unit – Pharmaceutical Sciences
 Vancouver General Hospital

Pharmacy Services




Conflicts of Interest

Research support	Nil
Employee	Nil
Consultant	Nil
Stakeholder	Nil
Speaker bureau	Nil
Scientific advisory board	Nil

Pharmacy Services



Ambulatory Leukemia-BMT Clinic



- 40-bed/chair clinic
- >14,000 visits last year
 - 113 AML & ALL patients for consolidation therapy
 - 64 auto-SCT for MM
 - 73 patients for allo-SCT supportive care until day +100
- Increase in patient referrals at a rate ~8%/yr

Pharmacy Services



Overview of Pharmacy Practice

- Facilitators of rational drug therapy
- Providers of pharmaceutical care
 - Monitor treatment plans
 - Educate patients and providers
 - Promote cost-effective therapy
- Identify patients at high risk for poor medication-related outcomes
 - All transplant patients
 - All patients receiving IV antimicrobials



Pharmacy Services



Outpatient Drug Therapy Challenges



- Delivery of outpatient IV antibiotics
- Obtaining drug benefit approval from Pharmacare
- Communication of medication information to outpatients

Pharmacy Services



Once Daily Vancomycin?

PHYSICIAN'S ORDERS

Leukemia/BMT Outpatient Parenteral Antibiotic Program

DATE: _____ TIME: _____

ONCE DAILY DOSING PARENTERAL ANTI-BIOTIC REGIMEN

1. Draw blood cultures and urine for CBC.
2. Check X-ray (if clinically applicable).
3. Patient fit for once daily dosing: No known "drug-drug" interactions.
4. Patient Wt _____ kg Ht _____ cm Ideal Wt _____ kg
5. Creatinine Wt _____ µmol/L

Calculations of Body Wt:
 Male: $(51.85 + 0.726 \times \text{cm} - 132.4) \div 4.545 = 58.076 \text{ kg}$
 Female: $(45.75 + 0.715 \times \text{cm} - 121.0) \div 4.545 = 54.027 \text{ kg}$

CRITICAL THINKING: Dosing Interval
 Change based on ideal body wt (IBW), except in obese patients (greater 120% of IBW) from one adjusted dose to:
 1. Tobramycin Q24H with creatinine clearance with CrCl₃₀ less than 30 mL/min
 2. Vancomycin Q24H with creatinine clearance with CrCl₃₀ less than 30 mL/min
 3. Less than or equal to 20 individualize

6. Patient check appropriate antibiotic:

TOBRAMYCIN _____ mg IV (6 mg/kg) IV Q _____ H according to the dosing interval table.

TOBRAMYCIN/VANCOMYCIN DOSING INTERVAL TABLE	CrCl (mL/min)	Dosing Interval (H)
Greater than or equal to 60	24	24
21 to 59	48	48
Less than or equal to 20	Individualize	

Estimated CrCl: $\frac{140 - \text{age}}{7.2} = 0.011 \times \text{SCr} (\mu\text{mol/L})$
 (x0.85 for females)

CEFTRIAXONE 2 g IV Q24H x 3 doses, then 1 g IV Q24H.

VANCOMYCIN _____ mg (20 mg/kg total body weight) IV Q _____ H over 90 minutes according to the dosing interval table. Patient to be reassessed in 72 hours for vancomycin discontinuation (if clinically improved/stable with no documented or suspected gram positive infection).

CEFTRIAXONE 2 g IV Q24H x 3 doses, then 1 g IV Q24H.

VANCOMYCIN _____ mg (20 mg/kg total body weight) IV Q _____ H over 90 minutes according to the dosing interval table. Patient to be reassessed in 72 hours for vancomycin discontinuation (if clinically improved/stable with no documented or suspected gram positive infection).

CEFTRIAXONE 2 g IV Q24H x 3 doses, then 1 g IV Q24H.

VANCOMYCIN _____ mg (20 mg/kg total body weight) IV Q _____ H over 90 minutes according to the dosing interval table. Patient to be reassessed in 72 hours for vancomycin discontinuation (if clinically improved/stable with no documented or suspected gram positive infection).

Physician Signature _____ Printed Name/PCP _____
 Date/Time _____

Pharmacy Services



Battling with PharmaCare

- Antifungal prophylaxis for transplant patients on steroids for GVHD
 - Fluconazole
 - Not a full benefit drug unless prescribed by BMT staff physician or PharmaCare Special Authority submitted
 - Voriconazole
 - Not a benefit drug unless PharmaCare Special Authority submitted for treatment of IA initiated in hospital
 - Posaconazole
 - Not a benefit drug, coverage considered for IA if intolerant or fail other therapies or for mucormycosis

Pharmacy Services



Are we effective communicators?

- 48 y/o male post-DCBT for ALL complicated by acute GVHD (gut) on prednisone
 - Steroid-induced DM on insulin NPH 17 units QAM

Pre-Supper Sugar <10 (<8)
↑ NPH dose by 2 units
every 2-3 days to target
Pre-Supper Sugar <10

Pharmacy Services



More Challenges Ahead...



"I don't think that's what the pharmacist meant when he said 'Take for two days and skip a day.'"

Pharmacy Services



Overcoming Challenges in Outpatient Drug therapy



Pharmacy Services

